

A28

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004006 A2

(51) International Patent Classification⁷: **A61K 31/00**

(21) International Application Number: **PCT/IB02/03295**

(22) International Filing Date: **28 June 2002 (28.06.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/301,407 **29 June 2001 (29.06.2001)** **US**

(71) Applicant (for all designated States except US): **AB SCIENCE [FR/FR]; 3, avenue George V, F-75008 Paris (FR).**

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MOUSSY, Alain [FR/FR]; 22 bis, passage Dauphine, F-75006 Paris (FR). KINET, Jean-Pierre [FR/US]; 3 Hunt Road, Lexington, MA 02421 (US).**

(74) Agents: **MARTIN, Jean-Jacques et al.; Cabinet Régimbeau, 20, rue de Chazelles, F-75847 Paris Cedex 17 (FR).**

(81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.**

(84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).**

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW**, **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)**
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **USE OF POTENT, SELECTIVE AND NON TOXIC C-KIT INHIBITORS FOR TREATING TUMOR ANGIOGENESIS**

(57) Abstract: The present invention relates to a method for inhibiting tumor angiogenesis comprising administering a c-kit inhibitor to a human in need of such treatment, more particularly a non toxic, potent and selective c-kit inhibitor, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

WO 03/004006 A2

WO 03/004006

PCT/IB02/03295

I

**Use of potent, selective and non toxic c-kit inhibitors for treating
tumor angiogenesis**

- 5 The present invention relates to a method for inhibiting tumor angiogenesis comprising administering a c-kit inhibitor to a human in need of such treatment, more particularly a non toxic, potent and selective c-kit inhibitor, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.
- 10 In 1971, Folkman J. (Tumor angiogenesis: Therapeutic implications, N. Engl. Jour. Med. 285:1182-1186) postulated that every increase in tumor cell population must be preceded by an increase in new capillaries converging on the tumor. Since, many evidence have accumulated demonstrating that the growth of new blood vessels from a preexisting microvascular bed is necessary for the growth, maintenance, and metastasis
- 15 of solid tumors.

Different compounds are being tried out for their potential therapeutic application in tumor angiogenesis. Among these compounds, Marimastat (British Biotech) and BMS-275291 (Bristol-Myers Squibb) are synthetic inhibitors of matrix metalloproteinases (MMPs), Neovastat (Aeterna) is a naturally occurring MMP inhibitor, Squalamine

20 (Magainin Pharmaceuticals) is extracted from dogfish shark liver, Endostatin (EntreMed) is an inhibitor of endothelial cells growth, SU5416 and SU6668 (Sugen) block VEGF / PDGF receptor signaling.

- 25 While these compounds block a particular stimulus leading to angiogenesis, they don't abolish all the pathways involved in the induction of new blood vessels, which results from the concomitant action of several growth factors and cytokines. These signals

WO 03/004006

PCT/IB02/03295

2

leading to tumor angiogenesis depend on the interaction of different tumor components : tumor parenchymal cells, endothelial cells, infiltrating cells from the bloodstream, and mast cells.

- 5 In connection with the present invention, we have determined that mast cells are in fact a major player in tumor angiogenesis due to their ability to secrete numerous growth factors and cytokines that ultimately balance the equilibrium in favor of vascular endothelial cells growth.
- 10 Mast cells (MC) are tissue elements derived from a particular subset of hematopoietic stem cells that express CD34, c-kit and CD13 antigens (Kirshenbaum et al, Blood. 94: 2333-2342, 1999 and Ishizaka et al, Curr Opin Immunol. 5: 937-43, 1993). Immature MC progenitors circulate in the bloodstream and differentiate in tissues. These differentiation and proliferation processes are under the influence of cytokines, one of
- 15 utmost importance being Stem Cell Factor (SCF), also termed Kit ligand (KL), Steel factor (SL) or Mast Cell Growth Factor (MCGF). SCF receptor is encoded by the protooncogene c-kit, that belongs to type III receptor tyrosine kinase subfamily (Boissan and Arock, J Leukoc Biol. 67: 135-48, 2000). This receptor is also expressed on others hematopoietic or non hematopoietic cells. Ligation of c-kit receptor by SCF induces its
- 20 dimerization followed by its transphosphorylation, leading to the recrutement and activation of various intracytoplasmic substrates. These activated substrates induce multiple intracellular signaling pathways responsible for cell proliferation and activation (Boissan and Arock, 2000). Mast cells are characterized by their heterogeneity, not only regarding tissue location and structure but also at the functional and histochemical levels
- 25 (Aldenborg and Enerback., Histochem. J. 26: 587-96, 1994 ; Bradding et al. J Immunol. 155: 297-307, 1995 ; Irani et al, J Immunol. 147: 247-53, 1991 ; Miller et al, Curr Opin Immunol. 1: 637-42, 1989 and Welle et al, J Leukoc Biol. 61: 233-45, 1997).

WO 03/004006

PCT/IB02/03295

3

Several observations have suggested the implication of mast cells in the pathogenesis of cancer and angiogenesis. First, mast cells have been shown to accumulate within and around solid tumors (Fisher E. and Fisher B. Role of mast cells in tumor growth. Arch. Pathol., 79: 185-191, 1965). Second, mast cells are distributed along blood vessels (Eady R. et al, Mast cell population density, blood vessel density and histamine content in normal skin. Br. J. Dermatol., 100: 635-640, 1979). Mast cell degranulation induces neovascularization in rat mesentery (Norrby K. et al, Mast-cell-mediated angiogenesis: a novel experimental model using the rat mesentery. Virchows Arch. B Cell Pathol. Incl. Mol. Pathol., 52: 195-206, 1986) and in the chick chorioallantoic membrane (Clinton M. et al. Effect of the mast cell activator compound 48/80 and heparin on angiogenesis in the chick chorioallantoic membrane. Int. J. Microcirc. Clin. Exp., 7: 315-326, 1988.).

Furthermore, when tumor cells are injected into a chick embryo, a 40-fold increase in mast cell density has been observed around the tumor implantation site compared with normal tissue (Kessler D. and Folkman J. Mast cells and tumor angiogenesis. Int. J. Cancer, 18: 703-709, 1976.). Injection of mast cell suspensions into animals induce an acceleration of tumor growth (Roche W., The nature and significance of tumor-associated mast cells. J. Pathol., 148: 175-182, 1986.), whereas decreasing the number of tissue mast cells leads to depression of tumor growth (Scott K., The mast cell, its amines, and tumor growth in rodents and man. Ann. NY Acad. Sci., 103: 285-312, 1963).

In addition, inhibiting mast cell degranulation with disodium cromoglycate has been demonstrated to significantly depress tumor growth (Ionov I., Inhibition of mast cell activity as a new approach to anticancer therapy. Int. J. Radiat. Biol., 60: 287-291, 1991). More recently, it has been suggested that mast cells in tumors modulate the

WO 03/004006

PCT/IB02/03295

4

neovascularization process (Wei Zhang et al , Modulation of Tumor Angiogenesis by Stem Cell Factor, Cancer Research 60, 6757-6762, December 1, 2000).

The present invention goes further based in the fact that tumor cell lines express stem
5 cell factor SCF and display c-kit receptors (Turner et al , Blood Volume 80, Issue 2, pp.
374-381, 1992). It is proposed here that tumor cells activate mast cells proliferation via
SCF, which in turn degranulate and release mediators such as histamine, TNF, IL-8,
VEGF or bFGF that acts together to promote angiogenesis. While blood vessels develop,
tumor is allowed to grow bigger, which results in an increase of SCF release.
10 Consequently, an activating feedback loop is created ultimately leading to further
activation of mast cells as well as growth of tumors and metastasis.

In addition, the role of mast cells in the process of tumor angiogenesis was confirmed by
comparing the rates of tumor vascularization, growth and metastasis in control
15 WBB6F1(-)/+ mice and in their mast-cell- deficient WBB6F1-W/W^v littermates
injected with MB49 murine bladder carcinoma cells. The results of these experiments
demonstrated that in mast-cell-deficient mice injected with tumor cells, there is a
decreased number of capillaries at the tumor periphery, reduced tumor size relative to
control mice, and an absence of metastases. These results have also shown that the
20 reduction of blood vessels at the tumor periphery might lead to a reduction in the number
of metastatic cells in mast-cell- deficient mice.

The relevance of the above mentioned hypothesis towards the human situation has been
confirmed by studies conducted in patients suffering from lung cancer, in whom mast
25 cell counts were significantly higher than in control normal tissues. Good correlation
was observed between intratumoral mast cell counts and microvessel counts. Double
staining showed highly angiogenic areas densely populated with mast cells. Importantly,

WO 03/004006

PCT/IB02/03295

5

members in the high mast cell count group had significantly worse prognosis than those in the low mast cell count group.

From these studies confirming a mutual activation between tumor cells and mast cells,
5 we can conclude that tumor-released vascular endothelial growth factors is related to mast cell accumulation, that intratumoral mast cells produce angiogenic factors, and that stromal mast cells correlate with angiogenesis and poor outcome in lung cancer.

In this regard, the general aim of the invention is to provide therapeutic strategies aiming
10 at blocking the activation and the survival of mast cells which are involved in tumor angiogenesis. This can be done by any means leading to mast cells death or inactivation. For example, it has been found that targeting c-kit or c-kit signaling is particularly suited to reach this goal. To this end, tyrosine kinase inhibitors that are non toxic and specific for mast cells are contemplated. These inhibitors are unable to promote death of IL-3
15 dependent cells cultured in presence of IL-3. Among such inhibitors, c-kit specific kinase inhibitors to inhibit mast cell proliferation, survival and activation are of a particular interest for clinical uses.

Description

20

Therefore, the present invention relates to a method for treating tumor angiogenesis comprising administering a tyrosine kinase inhibitor to a mammalian in need of such treatment, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

25

Tyrosine kinase inhibitors are selected for example from bis monocyclic, bicyclic or heterocyclic aryl compounds (WO 92/20642), vinylene-azaindole derivatives (WO 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones (US 5,330,992), Styryl compounds

WO 03/004006

PCT/IB02/03295

6

(US 5,217,999), styryl-substituted pyridyl compounds (US 5,302,606), seleoindoles and selenides (WO 94/03427), tricyclic polyhydroxylic compounds (WO 92/21660) and benzylphosphonic acid compounds (WO 91/15495), pyrimidine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones
5 (US 5,792,783, EP 934 931, US 5,834,504, US 5,883,116, US 5,883,113, US 5, 886,020, WO 96/40116 and WO 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520 722, US 3,772,295 and US 4,343,940) and aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO 95/15758).

10


Preferably, said tyrosine kinase inhibitor is a non-toxic, selective and potent c-kit inhibitor. Such inhibitors can be selected from pyrimidine derivatives such as N-phenyl-2-pyrimidine-amine derivatives (US 5,521,184 and WO 99/03854), indolinone derivatives and pyrrol-substituted indolinones (US 5,792,783, EP 934 931, US
15 5,834,504), US 5,883,116, US 5,883,113, US 5, 886,020, WO 96/40116 and WO 00/38519), as well as bis monocyclic, bicyclic aryl and heteroaryl compounds (EP 584 222, US 5,656,643 and WO 92/20642), quinazoline derivatives (EP 602 851, EP 520 722, US 3,772,295 and US 4,343,940), 4-amino-substituted quinazolines (US 3,470,182), 4-thienyl-2-(1H)-quinazolones, 6,7-dialkoxyquinazolines (US 3,800,039),
20 aryl and heteroaryl quinazoline (US 5,721,237, US 5,714,493, US 5,710,158 and WO 95/15758), 4-anilinoquinazoline compounds (US 4,464,375), and 4-thienyl-2-(1H)-quinazolones (US 3,551,427).

So, preferably, the invention relates to a method for preventing or treating tumor
25 angiogenesis comprising administering a non toxic, potent and selective c-kit inhibitor. Such inhibitor can be selected from pyrimidine derivatives, more particularly N-phenyl-2-pyrimidine-amine derivatives of formula I :

WO 03/004006


PCT/IB02/03295

7




wherein the R1, R2, R3, R13 to R17 groups have the meanings depicted in EP 564 409 B1, incorporated herein in the description.

- 5 Preferably, the N-phenyl-2-pyrimidine-amine derivative is selected from the compounds corresponding to formula II :



- 10 Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;
 R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;
 and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least
 15 one basic site, such as an amino function.

Preferably, R7 is the following group :



WO 03/004006

PCT/IB02/03295

8

Among these compounds, the preferred are defined as follows :

R1 is a heterocyclic group, especially a pyridyl group,


R2 and R3 are H,

5 R4 is a C1-C3 alkyl, especially a methyl group,

R5 and R6 are H,

and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one

basic site, such as an amino function, for example the group :




10

Therefore, in a preferred embodiment, the invention relates to a method for treating tumor angiogenesis comprising the administration of an effective amount of the compound known in the art as CGP57148B :

4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2

15 ylamino)phényl]-benzamide corresponding to the following formula :



The preparation of this compound is described in example 21 of EP 564 409 and the β -form, which is particularly useful is described in WO 99/03854.

20

WO 03/004006

PCT/IB02/03295

9

Alternatively, the invention relates to a method for treating tumor angiogenesis comprising administering a non toxic, potent and selective c-kit inhibitor to a mammalian in need of such treatment, selected from the group consisting of

- indolinone derivatives, more particularly pyrrol-substituted indolinones,
- 5 - monocyclic, bicyclic aryl and heteroaryl compounds, quinazoline derivatives,
- and quinaxolines, such as 2-phényl-quinaxoline derivatives, for example 2-phenyl-6,7-dimethoxy quinaxoline.

Preferably, said inhibitors are unable to promote death of IL-3 dependent cells cultured
10 in presence of IL-3.

In another embodiment, c-kit inhibitors as mentioned above are inhibitors of activated c-kit. In frame with the invention, the expression "activated c-kit" means a constitutively activated-mutant c-kit including at least one mutation selected from point mutations,
15 deletions, insertions, but also modifications and alterations of the natural c-kit sequence (SEQ ID N°1). Such mutations, deletions, insertions, modifications and alterations can occur in the transphosphorylase domain, in the juxtamembrane domain as well as in any domain directly or indirectly responsible for c-kit activity. The expression "activated c-kit" also means herein SCF-activated c-kit. Preferred and optimal SCF concentrations
20 for activating c-kit are comprised between 5.10^{-7} M and 5.10^{-6} M, preferably around 2.10^{-6} M. In a preferred embodiment, the activated-mutant c-kit in step a) has at least one mutation proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants. In another preferred embodiment, the activated-mutant c-kit in step a)
25 has a deletion in the juxtamembrane domain of c-kit. Such a deletion is for example between codon 573 and 579 called c-kit d(573-579). The point mutation V559G proximal to the juxtamembrane domain c-kit is also of interest.

WO 03/004006

PCT/IB02/03295

10

In this regard, the invention contemplates a method for treating tumor angiogenesis comprising administering to a mammalian in need of such treatment a compound that is a selective, potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :

- 5 a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested; under conditions allowing the components (i) and (ii) to form a complex,
b) selecting compounds that inhibit activated c-kit,
c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

10

This screening method can further comprise the step consisting of testing and selecting a subset of compounds identified in step b) that are inhibitors of mutant activated c-kit (for example in the transphosphorylase domain), which are also capable of inhibiting SCF-activated c-kit wild.

- 15 Alternatively, in step a) activated c-kit is SCF-activated c-kit wild.

A best mode for practicing this method consists of testing putative inhibitors at a concentration above 10 μ M in step a). Relevant concentrations are for example 10, 15, 20, 25, 30, 35 or 40 μ M.

20

In step c), IL-3 is preferably present in the culture media of IL-3 dependent cells at a concentration comprised between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

Examples of IL-3 dependent cells include but are not limited to :

- 25 - cell lines naturally expressing and depending on c-kit for growth and survival. Among such cells, human mast cell lines can be established using the following procedures :
normal human mast cells can be infected by retroviral vectors containing sequences coding for a mutant c-kit comprising the c-kit signal peptide and a TAG sequence

WO 03/004006

PCT/IB02/03295

11

allowing to differentiate mutant c-kits from c-kit wild expressed in hematopoietic cells by means of antibodies.

This technique is advantageous because it does not induce cellular mortality and the genetic transfer is stable and gives satisfactory yields (around 20 %). Pure normal human
5 mast cells can be routinely obtained by culturing precursor cells originating from blood obtained from human umbilical vein. In this regard, heparinated blood from umbilical vein is centrifuged on a Ficoll gradient so as to isolate mononucleated cells from other blood components. CD34+ precursor cells are then purified from the isolated cells mentioned above using the immunomagnetic selection system MACS (Miltenyi biotech).
10 CD34+ cells are then cultured at 37°C in 5 % CO₂ atmosphere at a concentration of 10⁵ cells per ml in the medium MCCM (α -MEM supplemented with L-glutamine, penicillin, streptomycin, 5 10⁻⁵ M β -mercaptoethanol, 20 % veal foetal serum, 1 % bovine albumin serum and 100 ng/ml recombinant human SCF. The medium is changed every 5 to 7 days. The percentage of mast cells present in the culture is assessed each week, using
15 May-Grünwal Giemsa or Toluidine blue coloration. Anti-tryptase antibodies can also be used to detect mast cells in culture. After 10 weeks of culture, a pure cellular population of mast cells (< 98 %) is obtained.

It is possible using standard procedures to prepare vectors expressing c-kit for transfecting the cell lines established as mentioned above. The cDNA of human c-kit has
20 been described in Yarden et al., (1987) EMBO J.6 (11), 3341-3351. The coding part of c-kit (3000 bp) can be amplified by PCR and cloned, using the following oligonucleotides :

- 5'AAGAAGAGATGGTACCTCGAGGGGTGACCC3' (SEQ ID No2) sens
- 5'CTGCTTCGCGGCCGCGTAACTCTTCTCAACCA3' (SEQ ID No3)

25 antisens

WO 03/004006

PCT/IB02/03295

12

The PCR products, digested with NotI and XhoI, has been inserted using T4 ligase in the pFlag-CMV vector (SIGMA), which vector is digested with NotI and XhoI and dephosphorylated using CIP (Biolabs). The pFlag-CMV-c-kit is used to transform bacterial clone XL1-blue. The transformation of clones is verified using the following

5 primers :

- 5'AGCTCGTTTAGTGAACCGTC3' (SEQ ID No4) sens,
- 5'GTCAGACAAAATGATGCAAC3' (SEQ ID No5) antisens.

Directed mutagenesis is performed using relevant cassettes is performed with routine and common procedure known in the art..

- 10 The vector Migr-1 (ABC) can be used as a basis for constructing retroviral vectors used for transfecting mature mast cells. This vector is advantageous because it contains the sequence coding for GFP at the 3' and of an IRES. These features allow to select cells infected by the retrovirus using direct analysis with a fluorocytometer. As mentioned above, the N-terminal sequence of c-kit c-DNA can be modified so as to introduce a Flag
- 15 sequence that will be useful to discriminating heterogeneous from endogenous c-kit:

Other IL-3 dependent cell lines that can be used include but are not limited to:

- BaF3 mouse cells expressing wild-type or mutated form of c-kit (in the juxtamembrane and in the catalytic sites) are described in Kitayama et al, (1996), Blood
- 20 88, 995-1004 and Tsujimura et al, (1999), Blood 93, 1319-1329.
- IC-2 mouse cells expressing either c-kit^{WT} or c-kit^{D814Y} are presented in Piao et al, (1996), Proc. Natl. Acad. Sci. USA 93, 14665-14669.

IL-3 independent cell lines are :

WO 03/004006

PCT/IB02/03295

13

- HMC-I, a factor-independent cell line derived from a patient with mast cell leukemia, expresses a juxtamembrane mutant c-kit polypeptide that has constitutive kinase activity (Furitsu T et al, J Clin Invest. 1993;92:1736-1744 ; Butterfield et al, Establishment of an immature mast cell line from a patient with mast cell leukemia. Leuk Res. 1988;12:345-355 and Nagata et al, Proc Natl Acad Sci U S A. 1995;92:10560-10564).
- P815 cell line (mastocytoma naturally expressing c-kit mutation at the 814 position) has been described in Tsujimura et al, (1994), Blood 83, 2619-2626.

The extent to which component (ii) inhibits activated c-kit can be measured *in vitro* or *in vivo*. In case it is measured *in vivo*, cell lines expressing an activated-mutant c-kit, which has at least one mutation proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants, are preferred.

Example of cell lines expressing an activated-mutant c-kit are as mentioned above.

15

In another preferred embodiment, the method further comprises the step consisting of testing and selecting compounds capable of inhibiting c-kit wild at concentration below 1 μ M. This can be measured *in vitro* or *in vivo*.

Therefore, compounds are identified and selected according to the method described above are potent, selective and non-toxic c-kit wild inhibitors.

20

Alternatively, the screening method according to the invention can be practiced *in vitro*. In this regard, the inhibition of mutant-activated c-kit and/or c-kit wild can be measured using standard biochemical techniques such as immunoprecipitation and western blot.

Preferably, the amount of c-kit phosphorylation is measured.

25

WO 03/004006

PCT/IB02/03295

14

In a still further embodiment, the invention contemplates a method for treating tumor angiogenesis as depicted above wherein the screening comprises :

- a) performing a proliferation assay with cells expressing a mutant c-kit (for example in the transphosphorylase domain), which mutant is a permanent activated c-kit, with a plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an $IC_{50} < 10 \mu M$, by measuring the extent of cell death,
- b) performing a proliferation assay with cells expressing c-kit wild said subset of candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-kit,
- c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting c-kit wild, each having an $IC_{50} < 10 \mu M$, preferably an $IC_{50} < 1 \mu M$, by measuring the extent of cell death.

Here, the extent of cell death can be measured by 3H thymidine incorporation, the trypan blue exclusion method or flow cytometry with propidium iodide. These are common techniques routinely practiced in the art.

Therefore, the invention embraces the use of the compounds defined above to manufacture a medicament for treating tumor angiogenesis in human.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries

WO 03/004006

PCT/IB02/03295

15

which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, Pa.).

5

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

10

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

20

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

25

WO 03/004006

PCT/IB02/03295

16

Pharmaceutical preparations which can be used orally include capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, 5 optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in 10 aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable 15 lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

20

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succine, acids, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred 25 preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

WO 03/004006

PCT/IB02/03295

17

Pharmaceutical compositions suitable for use in the invention include compositions wherein c-kit inhibitors are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. As mentioned above, a tyrosine kinase inhibitor and more particularly a c-kit inhibitor according to the invention is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

Therefore, the invention is also aimed at the use of a non toxic, potent and selective c-kit inhibitor for preparing a medicament for treating tumor angiogenesis in human, more particularly the use of a tyrosine kinase inhibitor or a c-kit inhibitor as defined above as being unable to promote death of IL-3 dependent cells cultured in presence of IL-3 for the manufacture of a medicament for treating tumor angiogenesis.

20

Utility of the invention will further ensue from the detailed description below.

Example 1: Identification of pro-angiogenic genes over-expressed in mast cells.

Genes expressed in mast cells, which contribute to the pathogenesis of diseases have been searched for. The purpose was to identify 1) genes expressed in different type of mast cells involved in different forms of mastocytosis and caused by mutations on the c-kit receptor and 2) genes expressed in mast cells involved in different pathologies,

WO 03/004006

PCT/IB02/03295

18

especially in the development of solid tumors, in metatasis as well as in inflammatory syndromes.

In a first approach, genes whose expression is linked to the differentiation of mast cells
 5 were identified from totipotent CD34+ cells, immature hematopoietic cells in course of differentiation and normal mature mast cells.

partial cDNA expression arrays

10 Expression profile of CD34+ cells extracted from human bone marrow and expression profile of mature mast cells derived from these CD 34+ induced by Stem Cell Factor (SCF) were obtained and analyzed by the Atlas Software.

Two types of membranes were used. The first one allows to detect 588 genes which are
 15 « general » and the other one allows to detect genes belonging to the haematology domain.

Genes whose expression is significantly increased ($\geq \times 3$) during mast cells differentiation are shown in the Table I below:

20

TABLE I: PARTIAL TRANSCRIPTOME OF MAST CELLS

Membrane « general »

			Genes over-expressed in mast cells versus CD34+ cells
code	Ratio	Diff	Protein/gene
F3j	4.008417	26450	transcription factor ETR103; early growth response protein 1 (EGR-1) (KROX24); zinc finger protein 225 (AT225)
C3i	4.694429	27590	Notch4
E2d	5.850825	16454	TIMP-3; mitogen-inducible gene 5 (mig-5)
C3j	6.289161	16005	Jagged 1

WO 03/004006

PCT/IB02/03295

19

F6e	6.525074	14984	platelet-derived growth factor A chain (PDGF-A)
B6b	7.194957	30219	C-kit
B6c	7.476532	26217	proto-oncogene c-src l tyrosine kinase domain
E1f	9.513902	18986	MMP-9; gelatinase B
D6h	10.915376	16638	LAR
B6a	11.612540	13202	C-fos
E1n	23.914141	13611	MMP-17 (MT4-MMP)

Membrane « haematology »

5

Up regulated genes cut <3	
Monocyte chemotactic factor	38,9
IL1 receptor antagonist	33,9
DNA binding protein inhibitor	27,5
CD9 antigen; p24; MIC3	11,3
RGS1 B-cell activation prot	11,7
LIF; differentiation-stimulating factor	10,1
ICAM1; CD 54 antigen	9,4
ST2 protein precursor	7,9
GATA2	4,7
BTK	4,1
JAK3	3,8
CD44 precursor	3,5

Over-expression Notch4 and Jagged 1

10

Differentiation of CD 43+ in mast cells results in a concomitant increase of the expression of **Notch4** and its ligand **Jagged 1**.

- 15 Notch 4 is a membrane receptor present in embryonic cells and in the endothelium. Jagged and notch4 are involved in the mechanism leading to angiogenesis. Notch signaling can regulate the angiogenic process since Notch4/int-3 and Jagged-1 are able to induce cultured endothelial cells to form cellular structures with morphological and

WO 03/004006

PCT/IB02/03295

20

biochemical properties of endothelial microvessels; Uyttendaele H. et al, Notch4/int-3, a mammary proto-oncogene, is an endothelial cell-specific Notch gene, Development. 122: 2251-59 (1996) and Uyttendaele, H. et al (2000) Notch4 and Jagged1 induce microvessel differentiation of rat brain endothelial cells. Microvascular Res.

5 Volkhard L. et al, (Am J Pathol 2001, 159:875–883) also reported that Jagged regulation of cell-cell and cell-matrix interactions may contribute to the control of cell migration in situations of tissue remodeling *in vivo*.

In conclusion, secreted Jagged 1 can act at the level of vascular endothelium (cells
10 expressing notch4) and induce the vascularization mechanism.

The autocrine and paracrine Jagged / Notch4 system in mast cells can contribute to angiogenesis. These results demonstrate that mast cells are effector cells of angiogenesis.

15

20

WO 03/004006

PCT/IB02/03295

21

CLAIMS


1. A method for treating tumor angiogenesis comprising administering a tyrosine kinase inhibitor to a mammalian in need of such treatment, wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.
2. A method according to claim 1, wherein said tyrosine kinase inhibitor is a non-toxic, selective and potent c-kit inhibitor.
3. A method according to claim 2, wherein said inhibitor is selected from the group consisting of indolinones, pyrimidine derivatives, pyrrolopyrimidine derivatives, quinazoline derivatives, quinoxaline derivatives, pyrazoles derivatives, bis monocyclic, bicyclic or heterocyclic aryl compounds, vinylene-azaindole derivatives and pyridyl-quinolones derivatives, styryl compounds, styryl-substituted pyridyl compounds, , seleoindoles, selenides, tricyclic polyhydroxylic compounds and benzylphosphonic acid compounds.
4. A method for treating tumor angiogenesis comprising administering a non toxic, potent and selective c-kit inhibitor to a mammalian in need of such treatment, selected from the group consisting of :
- pyrimidine derivatives, more particularly N-phenyl-2-pyrimidine-amine derivatives.
 - indolinone derivatives, more particularly pyrrol-substituted indolinones,
 - monocyclic, bicyclic aryl and heteroaryl compounds,
 - and quinazoline derivatives,
- wherein said inhibitor is unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

WO 03/004006

PCT/IB02/03295

22

5. A method according to claim 4, wherein said inhibitor is a N-phenyl-2-pyrimidine-amine derivative selected from the compounds corresponding to formula II :



5


Wherein R1, R2 and R3 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl or a cyclic or heterocyclic group, especially a pyridyl group;

R4, R5 and R6 are independently chosen from H, F, Cl, Br, I, a C1-C5 alkyl, especially a methyl group;

10 and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function.

6. A method according to claim 5, wherein R1 is a heterocyclic group, especially a pyridyl group, R2 and R3 are H, R4 is a C1-C3 alkyl, especially a methyl group, R5 and
15 R6 are H, and R7 is a phenyl group bearing at least one substituent, which in turn possesses at least one basic site, such as an amino function.

7. A method according to claim 6, wherein R7 is the following group :



20

8. A method according to claim 4, wherein said inhibitor is the 4-(4-méthylpipérazine-1-ylméthyl)-N-[4-méthyl-3-(4-pyridine-3-yl)pyrimidine-2 ylamino]phényl]-benzamide.

WO 03/004006

PCT/IB02/03295

23

9. A method according to claim 4, wherein said inhibitor is an inhibitor of activated c-kit selected from a constitutively activated-mutant c-kit and/or SCF-activated c-kit.

5 10. A method according to claim 9, wherein the activated-mutant c-kit has at least one mutation selected from mutations proximal to Y823, more particularly between amino acids 800 to 850 of SEQ ID No1 involved in c-kit autophosphorylation, notably the D816V, D816Y, D816F and D820G mutants, and a deletion in the juxtamembrane domain of c-kit, preferably between codon 573 and 579.

10

11. A method for treating tumor angiogenesis comprising administering to a mammalian in need of such treatment a compound that is a selective, potent and non toxic inhibitor of activated c-kit obtainable by a screening method which comprises :

a) bringing into contact (i) activated c-kit and (ii) at least one compound to be tested;
15 under conditions allowing the components (i) and (ii) to form a complex,
b) selecting compounds that inhibit activated c-kit,
c) testing and selecting a subset of compounds identified in step b), which are unable to promote death of IL-3 dependent cells cultured in presence of IL-3.

20 12. A method according to claim 11, wherein the screening method further comprises the step consisting of testing and selecting a subset of compounds identified in step b) that are inhibitors of mutant activated c-kit, which are also capable of inhibiting SCF-activated c-kit wild.

25 13. A method according to claim 11, wherein activated c-kit is SCF-activated c-kit wild.

14. A method according to one of claims 11 to 13, wherein putative inhibitors are tested at a concentration above 10 μ M in step a).

WO 03/004006

PCT/IB02/03295

24

15. A method according to one of claims 11 to 14, wherein IL-3 is present in the culture media of IL-3 dependent cells at a concentration comprised between 0.5 and 10 ng/ml, preferably between 1 to 5 ng/ml.

5 16. A method according to one of claims 11 to 15, wherein the extent to which component (ii) inhibits activated c-kit can be measured *in vitro* or *in vivo*.

17. A method according to one of claims 11 to 16 wherein, the screening method further comprises the step consisting of testing and selecting *in vitro* or *in vivo* compounds
10 capable of inhibiting c-kit wild at concentration below 1 μ M.

18. A method according to claim 17 wherein, wherein the test is performed using cells lines selected from the group consisting of mast cells, transfected mast cells, BaF3, and IC-2.
15

19. A method according to claim 17, wherein the test includes the determination of the amount of c-kit phosphorylation.

20. A method for treating tumor angiogenesis according to one of claims 11 to 18, wherein the screening comprises :
20

- a) performing a proliferation assay with cells expressing a mutant c-kit (for example in the transphosphorylase domain), which mutant is a permanent activated c-kit, with a plurality of test compounds to identify a subset of candidate compounds targeting activated c-kit, each having an $IC_{50} < 10 \mu M$, by measuring the extent of cell death,
- 25 b) performing a proliferation assay with cells expressing c-kit wild said subset of candidate compounds identified in step (a), said cells being IL-3 dependent cells cultured in presence of IL-3, to identify a subset of candidate compounds targeting specifically c-kit,

WO 03/004006

PCT/IB02/03295

25

c) performing a proliferation assay with cells expressing c-kit, with the subset of compounds identified in step b) and selecting a subset of candidate compounds targeting c-kit wild, each having an $IC_{50} < 10 \mu M$, preferably an $IC_{50} < 1 \mu M$, by measuring the extent of cell death.

5

21. A method according to one of claims 1 to 20 for treating tumor angiogenesis in human.

22. A method according to claim 21, wherein the inhibitor is administered orally.

10

23. A method according to claim 21, wherein the inhibitor is administered topically

24. Use of a non toxic, potent and selective c-kit inhibitor for preparing a medicament for treating tumor angiogenesis in human.

15

WO 03/004006

1

PCT/IB02/03295

SEQUENCE LISTING

<110> AB Science

<120> Use of potent, selective and non toxic c-kit inhibitors for treating tumor angiogenesis

<130> D19702 NT

<150> US 60/301,407

<151> 2001-06-29

<160> 5

<170> PatentIn Ver. 2.1

<210> 1

<211> 976

<212> PRT

<213> Homo sapiens

<220>

<223> Human c-kit

<400> 1

Met Arg Gly Ala Arg Gly Ala Trp Asp Phe Leu Cys Val Leu Leu Leu
 1 5 10 15

Leu Leu Arg Val Gln Thr Gly Ser Ser Gln Pro Ser Val Ser Pro Gly
 20 25 30

Glu Pro Ser Pro Pro Ser Ile His Pro Gly Lys Ser Asp Leu Ile Val
 35 40 45

Arg Val Gly Asp Glu Ile Arg Leu Leu Cys Thr Asp Pro Gly Phe Val
 50 55 60

Lys Trp Thr Phe Glu Ile Leu Asp Glu Thr Asn Glu Asn Lys Gln Asn
 65 70 75 80

Glu Trp Ile Thr Glu Lys Ala Glu Ala Thr Asn Thr Gly Lys Tyr Thr
 85 90 95

Cys Thr Asn Lys His Gly Leu Ser Asn Ser Ile Tyr Val Phe Val Arg
 100 105 110

Asp Pro Ala Lys Leu Phe Leu Val Asp Arg Ser Leu Tyr Gly Lys Glu
 115 120 125

Asp Asn Asp Thr Leu Val Arg Cys Pro Leu Thr Asp Pro Glu Val Thr
 130 135 140

Asn Tyr Ser Leu Lys Gly Cys Gln Gly Lys Pro Leu Pro Lys Asp Leu
 145 150 155 160

Arg Phe Ile Pro Asp Pro Lys Ala Gly Ile Met Ile Lys Ser Val Lys
 165 170 175

Arg Ala Tyr His Arg Leu Cys Leu His Cys Ser Val Asp Gln Glu Gly
 180 185 190

WO 03/004006

2

PCT/IB02/03295

Lys Ser Val Leu Ser Glu Lys Phe Ile Leu Lys Val Arg Pro Ala Phe
 195 200 205
 Lys Ala Val Pro Val Val Ser Val Ser Lys Ala Ser Tyr Leu Leu Arg
 210 215 220
 Glu Gly Glu Glu Phe Thr Val Thr Cys Thr Ile Lys Asp Val Ser Ser
 225 230 235 240
 Ser Val Tyr Ser Thr Trp Lys Arg Glu Asn Ser Gln Thr Lys Leu Gln
 245 250 255
 Glu Lys Tyr Asn Ser Trp His His Gly Asp Phe Asn Tyr Glu Arg Gln
 260 265 270
 Ala Thr Leu Thr Ile Ser Ser Ala Arg Val Asn Asp Ser Gly Val Phe
 275 280 285
 Met Cys Tyr Ala Asn Asn Thr Phe Gly Ser Ala Asn Val Thr Thr Thr
 290 295 300
 Leu Glu Val Val Asp Lys Gly Phe Ile Asn Ile Phe Pro Met Ile Asn
 305 310 315 320
 Thr Thr Val Phe Val Asn Asp Gly Glu Asn Val Asp Leu Ile Val Glu
 325 330 335
 Tyr Glu Ala Phe Pro Lys Pro Glu His Gln Gln Trp Ile Tyr Met Asn
 340 345 350
 Arg Thr Phe Thr Asp Lys Trp Glu Asp Tyr Pro Lys Ser Glu Asn Glu
 355 360 365
 Ser Asn Ile Arg Tyr Val Ser Glu Leu His Leu Thr Arg Leu Lys Gly
 370 375 380
 Thr Glu Gly Gly Thr Tyr Thr Phe Leu Val Ser Asn Ser Asp Val Asn
 385 390 395 400
 Ala Ala Ile Ala Phe Asn Val Tyr Val Asn Thr Lys Pro Glu Ile Leu
 405 410 415
 Thr Tyr Asp Arg Leu Val Asn Gly Met Leu Gln Cys Val Ala Ala Gly
 420 425 430
 Phe Pro Glu Pro Thr Ile Asp Trp Tyr Phe Cys Pro Gly Thr Glu Gln
 435 440 445
 Arg Cys Ser Ala Ser Val Leu Pro Val Asp Val Gln Thr Leu Asn Ser
 450 455 460
 Ser Gly Pro Pro Phe Gly Lys Leu Val Val Gln Ser Ser Ile Asp Ser
 465 470 475 480
 Ser Ala Phe Lys His Asn Gly Thr Val Glu Cys Lys Ala Tyr Asn Asp
 485 490 495
 Val Gly Lys Thr Ser Ala Tyr Phe Asn Phe Ala Phe Lys Gly Asn Asn
 500 505 510
 Lys Glu Gln Ile His Pro His Thr Leu Phe Thr Pro Leu Leu Ile Gly

WO 03/004006

3

PCT/IB02/03295

515					520					525					
Phe	Val	Ile	Val	Ala	Gly	Met	Met	Cys	Ile	Ile	Val	Met	Ile	Leu	Thr
530						535					540				
Tyr	Lys	Tyr	Leu	Gln	Lys	Pro	Met	Tyr	Glu	Val	Gln	Trp	Lys	Val	Val
545					550					555					560
Glu	Glu	Ile	Asn	Gly	Asn	Asn	Tyr	Val	Tyr	Ile	Asp	Pro	Thr	Gln	Leu
				565					570					575	
Pro	Tyr	Asp	His	Lys	Trp	Glu	Phe	Pro	Arg	Asn	Arg	Leu	Ser	Phe	Gly
			580					585					590		
Lys	Thr	Leu	Gly	Ala	Gly	Ala	Phe	Gly	Lys	Val	Val	Glu	Ala	Thr	Ala
		595					600					605			
Tyr	Gly	Leu	Ile	Lys	Ser	Asp	Ala	Ala	Met	Thr	Val	Ala	Val	Lys	Met
	610					615					620				
Leu	Lys	Pro	Ser	Ala	His	Leu	Thr	Glu	Arg	Glu	Ala	Leu	Met	Ser	Glu
						630					635				640
Leu	Lys	Val	Leu	Ser	Tyr	Leu	Gly	Asn	His	Met	Asn	Ile	Val	Asn	Leu
				645					650					655	
Leu	Gly	Ala	Cys	Thr	Ile	Gly	Gly	Pro	Thr	Leu	Val	Ile	Thr	Glu	Tyr
			660					665					670		
Cys	Cys	Tyr	Gly	Asp	Leu	Leu	Asn	Phe	Leu	Arg	Arg	Lys	Arg	Asp	Ser
		675					680					685			
Phe	Ile	Cys	Ser	Lys	Gln	Glu	Asp	His	Ala	Glu	Ala	Ala	Leu	Tyr	Lys
						695					700				
Asn	Leu	Leu	His	Ser	Lys	Glu	Ser	Ser	Cys	Ser	Asp	Ser	Thr	Asn	Glu
						710					715				720
Tyr	Met	Asp	Met	Lys	Pro	Gly	Val	Ser	Tyr	Val	Val	Pro	Thr	Lys	Ala
				725					730					735	
Asp	Lys	Arg	Arg	Ser	Val	Arg	Ile	Gly	Ser	Tyr	Ile	Glu	Arg	Asp	Val
			740					745					750		
Thr	Pro	Ala	Ile	Met	Glu	Asp	Asp	Glu	Leu	Ala	Leu	Asp	Leu	Glu	Asp
		755					760					765			
Leu	Leu	Ser	Phe	Ser	Tyr	Gln	Val	Ala	Lys	Gly	Met	Ala	Phe	Leu	Ala
						775					780				
Ser	Lys	Asn	Cys	Ile	His	Arg	Asp	Leu	Ala	Ala	Arg	Asn	Ile	Leu	Leu
						790					795				800
Thr	His	Gly	Arg	Ile	Thr	Lys	Ile	Cys	Asp	Phe	Gly	Leu	Ala	Arg	Asp
				805					810					815	
Ile	Lys	Asn	Asp	Ser	Asn	Tyr	Val	Val	Lys	Gly	Asn	Ala	Arg	Leu	Pro
			820					825					830		
Val	Lys	Trp	Met	Ala	Pro	Glu	Ser	Ile	Phe	Asn	Cys	Val	Tyr	Thr	Phe
		835					840					845			

WO 03/004006

4

PCT/IB02/03295

Glu Ser Asp Val Trp Ser Tyr Gly Ile Phe Leu Trp Glu Leu Phe Ser
 850 855 860
 Leu Gly Ser Ser Pro Tyr Pro Gly Met Pro Val Asp Ser Lys Phe Tyr
 865 870 875 880
 Lys Met Ile Lys Glu Gly Phe Arg Met Leu Ser Pro Glu His Ala Pro
 885 890 895
 Ala Glu Met Tyr Asp Ile Met Lys Thr Cys Trp Asp Ala Asp Pro Leu
 900 905 910
 Lys Arg Pro Thr Phe Lys Gln Ile Val Gln Leu Ile Glu Lys Gln Ile
 915 920 925
 Ser Glu Ser Thr Asn His Ile Tyr Ser Asn Leu Ala Asn Cys Ser Pro
 930 935 940
 Asn Arg Gln Lys Pro Val Val Asp His Ser Val Arg Ile Asn Ser Val
 945 950 955 960
 Gly Ser Thr Ala Ser Ser Ser Gln Pro Leu Leu Val His Asp Asp Val
 965 970 975

<210> 2
 <211> 30
 <212> DNA
 <213> Homo sapiens

<220>
 <223> Primer

<400> 2
 aagaagagat ggtacctcga ggggtgaccc

30

<210> 3
 <211> 33
 <212> DNA
 <213> Homo sapiens

<220>
 <223> Primer

<400> 3
 ctgcttcgcg gccgcgttaa ctcttctcaa cca

33

<210> 4
 <211> 20
 <212> DNA
 <213> Homo sapiens

<220>
 <223> Primer

WO 03/004006

5

PCT/IB02/03295

<400> 4

agctcgttta gtgaaccgtc

20

<210> 5

<211> 20

<212> DNA

<213> Homo sapiens

<220>

<223> Primer

<400> 5

gtcagacaaa atgatgcaac

20